

Direct Inhibition of the Longevity-Promoting Factor SKN-1 by Insulin-like Signaling in *C. elegans*

Jennifer M.A. Tullet,^{1,5} Maren Hertweck,^{2,5} Jae Hyung An,^{1,3,5} Joseph Baker,¹ Ji Yun Hwang,³ Shu Liu,² Riva P. Oliveira,¹ Ralf Baumeister,^{2,4} and T. Keith Blackwell^{1,*}

¹Section on Developmental and Stem Cell Biology, Joslin Diabetes Center; Department of Pathology, Harvard Medical School; Harvard Stem Cell Institute, 1 Joslin Place, Boston, MA 02215, USA

²Center of Biochemistry and Molecular Cell Research (ZBMZ, Faculty of Medicine), Bioinformatics and Molecular Genetics (Faculty of Biology), University of Freiburg, Schänzlestr. 1, D-79104 Freiburg, Germany

³Protein Network Research Center, Yonsei University, 134 Shinchon-dong, Seodaemun-gu, Seoul, 120-749, Republic of Korea

⁴Center for Systems Biology (ZBSA), University of Freiburg, D-79104 Freiburg, Germany

⁵These authors contributed equally to this work.

*Correspondence: keith.blackwell@joslin.harvard.edu

DOI 10.1016/j.cell.2008.01.030

SUMMARY

Insulin/IGF-1-like signaling (IIS) is central to growth and metabolism and has a conserved role in aging. In *C. elegans*, reductions in IIS increase stress resistance and longevity, effects that require the IIS-inhibited FOXO protein DAF-16. The *C. elegans* transcription factor SKN-1 also defends against oxidative stress by mobilizing the conserved phase 2 detoxification response. Here we show that IIS not only opposes DAF-16 but also directly inhibits SKN-1 in parallel. The IIS kinases AKT-1, -2, and SGK-1 phosphorylate SKN-1, and reduced IIS leads to constitutive SKN-1 nuclear accumulation in the intestine and SKN-1 target gene activation. SKN-1 contributes to the increased stress tolerance and longevity resulting from reduced IIS and delays aging when expressed transgenically. Furthermore, SKN-1 that is constitutively active increases life span independently of DAF-16. Our findings indicate that the transcription network regulated by SKN-1 promotes longevity and is an important direct target of IIS.

INTRODUCTION

In many studies of aging, extended longevity correlates with enhanced resistance against oxidative damage and other stresses (Finkel and Holbrook, 2000; Lithgow and Walker, 2002). In *C. elegans*, both life-span and stress resistance can be increased by the alteration of several processes, including insulin/IGF-1-like signaling (IIS) (Friedman and Johnson, 1988; Gottlieb and Ruvkun, 1994; Kenyon et al., 1993; Larsen, 1993), caloric intake (Houthoofd et al., 2002), mitochondrial respiration (Dillin et al., 2002; Lakowski and Hekimi, 1996; Lee et al., 2003), and germline function (Arantes-Oliveira et al., 2002). It is a formidable but crit-

ical challenge to understand how modulation of these processes might inhibit or defend against stresses that contribute to aging.

Reductions in IIS have been associated with increased stress resistance and longevity in diverse species (Kenyon, 2005). Studies in worms and flies have established the paradigm that these effects derive from increased activity of the FOXO transcription factors, which are inhibited through IIS-induced phosphorylation (Figure 1A) (Antebi, 2007; Kenyon, 2005). FOXO proteins are also important in many of the biological effects of insulin in mammals (Accilli and Arden, 2004; van der Horst and Burgering, 2007). In *C. elegans*, signaling through the IIS receptor DAF-2 ultimately directs the related kinases AKT-1, -2, and SGK-1 to phosphorylate the FOXO protein DAF-16, thereby inhibiting its accumulation in nuclei (Henderson and Johnson, 2001; Hertweck et al., 2004; Lin et al., 2001; Ogg et al., 1997). DAF-16 is required for phenotypes that are associated with decreased IIS, including increases in stress resistance, longevity, and the propensity to undergo diapause, the formation of long-lived dauer larvae that can survive adverse environmental conditions (Kenyon, 2005; Kenyon et al., 1993; Kimura et al., 1997). DAF-16 regulates genes that represent diverse processes, including resistance to oxidative and other stresses (Antebi, 2007; Kenyon, 2005).

The phase 2 detoxification response also provides a conserved defense against oxidative stress and consists of numerous enzymes that scavenge free radicals and other reactive compounds and produce or transfer glutathione (McMahon et al., 2001). In *C. elegans* the phase 2 response is orchestrated by the transcription factor SKN-1. SKN-1 initiates development of the feeding and digestive system during the earliest embryonic stages, then is required postembryonically for normal life span and stress resistance (An and Blackwell, 2003; Bowerman et al., 1992). SKN-1 is present in nuclei constitutively in the ASI neurons (putative hypothalamus) (An and Blackwell, 2003), where it is required for longevity to be extended by dietary restriction (DR), a condition that increases longevity in organisms as diverse as yeast and rodents (Bishop and Guarente, 2007). The stress resistance function of




Figure 1. Accumulation of SKN-1 in Intestinal Nuclei Is Inhibited by DAF-2

(A) A partial schematic of *C. elegans* insulin-like signaling and our working model. (B) *skn-1* isoforms, alleles, and transgenes. The SKN-1A isoform is transcribed from an upstream operon promoter and differs from SKN-1C by the addition of 90 amino acids at its N terminus. SKN-1B and SKN-1C are each expressed from their own distinct upstream promoters (An and Blackwell, 2003; Bishop and Guarente, 2007). The *skn-1(zu67)*, *(zu129)*, and *(zu135)* mutations each create a premature stop codon (B. Bowerman, personal communication). Green fluorescent protein (GFP) is fused to the C terminus of SKN-1 in the transgenes indicated at the bottom. Strains that carry these and other transgenes are described in Table S2. The *skn-1a/c* RNAi construct includes only coding sequence and targets both SKN-1A and SKN-1C but does not alter the levels of SKN-1B in the ASI neurons (Bishop and Guarente, 2007). (C) SKN-1::GFP accumulates in intestinal nuclei when DAF-2 activity is reduced. (D) Intestinal expression of *SKN-1B/C::GFP* in *daf-2(e1370)* derives from SKN-1C. (E) Quantification of SKN-1 intestinal nuclear accumulation. Here and in Figure 2, results of experiments performed on L4 larvae have been combined and scored as described in the Supplemental Experimental Procedures and in An and Blackwell (2003) and An et al. (2005). p values were derived from a χ^2 test.

SKN-1 is mediated by its expression in the intestine (digestive system) (Bishop and Guarente, 2007), where SKN-1 accumulates in nuclei and activates phase 2 gene expression inducibly in response to stress (An and Blackwell, 2003). In the intestine, phosphorylation of SKN-1 by p38/mitogen-activated protein kinase (MAPK) signaling is required for its accumulation in nuclei, while negative regulation via glycogen synthase kinase-3 (GSK-3) phosphorylation is needed to prevent this from occurring constitutively (Figure 1A) (An et al., 2005; Inoue et al., 2005).

As *daf-16* is required for the increased stress resistance and longevity that are seen when IIS is reduced, it has seemed likely that these effects of decreased IIS could be accounted for by

increased DAF-16 activity. However, we reasoned that if it is advantageous for IIS to inhibit stress response genes by acting on DAF-16, then IIS might also oppose SKN-1 (Figure 1A). Accordingly, here we show that reductions in IIS cause SKN-1 to accumulate constitutively in intestinal nuclei in the absence of stress and to activate phase 2 target genes. Importantly, these events do not require *daf-16*. AKT-1, -2, and SGK-1 phosphorylate SKN-1 at multiple sites, and mutation of an AKT phosphorylation site results in SKN-1 being present in intestinal nuclei constitutively. *skn-1* mutations significantly suppress the oxidative stress resistance and longevity phenotypes associated with reduced IIS. Aging is delayed when SKN-1 is expressed transgenically,

and a mutant SKN-1 form that localizes constitutively to intestinal nuclei increases life span in the absence of *daf-16*. In summary, our data indicate that IIS directly inhibits SKN-1 in parallel to DAF-16, that SKN-1 contributes to the stress and longevity phenotypes associated with reduced IIS, and that SKN-1 defines an important pro-longevity mechanism.

RESULTS

DAF-2 Signaling Inhibits SKN-1 Accumulation in Intestinal Nuclei

If IIS inhibits SKN-1 analogously to DAF-16 (Figure 1A), then reduced DAF-2 signaling should allow SKN-1 to accumulate in intestinal nuclei. To test this idea, we first investigated how reductions in DAF-2 activity affect the distribution of SKN-1 that is expressed from the previously described *SKN-1B/C::GFP* transgene (Figure 1B). This transgene rescues the maternal lethality, stress sensitivity, and DR-associated longevity defects of *skn-1* mutants (An and Blackwell, 2003; An et al., 2005; Bishop and Guarente, 2007). *SKN-1B/C::GFP* encodes two of three SKN-1 isoforms, SKN-1C and SKN-1B, which are expressed in the intestine and ASI neurons, respectively (Bishop and Guarente, 2007). In wild-type (WT) animals that carry extrachromosomal or integrated *SKN-1B/C::GFP*, SKN-1::GFP is generally undetectable in intestinal nuclei except under stress conditions or after inhibition of its phosphorylation by GSK-3 (An and Blackwell, 2003; An et al., 2005). In contrast, SKN-1::GFP was constitutively present in intestinal nuclei of transgenic *daf-2(e1370)* and *daf-2(e1368)* animals or in WT transgenics exposed to *daf-2* RNA interference (RNAi) at either 20°C or low temperature (16°C) (Figures 1C and 1E; see also Figures S1A–S1C available online). This nuclear SKN-1 appeared to correspond to SKN-1C because RNAi against this isoform prevented SKN-1::GFP from accumulating in intestinal nuclei in response to *daf-2* mutation (Figure 1D).

We also analyzed a transgene that encodes all three SKN-1 isoforms in the context of a large operon (*SKN-1op::GFP*; Figure 1B). In the WT background this transgene was expressed in a similar pattern to *SKN-1B/C::GFP* (not shown), and in *daf-2(e1370)* SKN-1 that was expressed from *SKN-1op::GFP* also accumulated in intestinal nuclei constitutively (Figure 1E). In contrast to these effects in the intestine, *daf-2* mutation did not obviously alter the levels of nuclear SKN-1::GFP in the ASI neurons (Figures 1C and 1D; data not shown). We conclude that signaling through DAF-2 may be required to prevent SKN-1 from accumulating constitutively in intestinal nuclei.

DAF-2 Signaling Inhibits SKN-1 Directly, through Phosphorylation

Signaling through DAF-2 ultimately activates the AKT-1, -2, and SGK-1 kinases, which phosphorylate and inhibit DAF-16 (Figure 1A) (Cahill et al., 2001; Hertweck et al., 2004; Lin et al., 2001). RNAi knockdown of *akt-1*, *akt-2* or *sgk-1* dramatically increased the presence in intestinal nuclei of SKN-1::GFP that was expressed from either *SKN-1B/C::GFP* or *SKN-1op::GFP* (Figures 2A and 2B), supporting the model that IIS inhibits SKN-1 directly. Importantly, *daf-16* was not required for SKN-1 expression, or for the nuclear accumulation of SKN-1 that re-

sulted from reductions in IIS. A constitutively nuclear SKN-1 substitution mutant in which a critical inhibitory GSK-3 phosphorylation site has been altered (SKN-1(S393A); An et al., 2005) was appropriately present in intestinal nuclei in null *daf-16* animals (Figure S2). In addition, in a null *daf-16* background, RNAi of the IIS kinases still increased the levels of intestinal nuclear SKN-1::GFP expressed from *SKN-1op::GFP* (Figure 2A). Finally, in *daf-2(e1370)* animals, the nuclear accumulation of SKN-1 expressed from *SKN-1B/C::GFP* was not reduced by *daf-16* RNAi (Figure S1D). We conclude that the IIS pathway inhibits SKN-1 in the intestine, and that this regulation occurs independently of DAF-16.

We previously observed that p38 MAPK signaling is required for SKN-1 to accumulate in intestinal nuclei not only in response to stress but also when GSK-3 phosphorylation of SKN-1 is inhibited under normal conditions (An et al., 2005; Inoue et al., 2005). In the latter case, even though negative regulation of SKN-1 is impaired, the low-level background p38 signaling that is seen under normal conditions is still required. To test whether p38 signaling is also needed for SKN-1 to be present in intestinal nuclei when IIS is reduced, we performed *akt-1*; *akt-2* and *sgk-1* RNAi in animals that lacked the p38 pathway MAPK kinase SEK-1 (Inoue et al., 2005). In *sek-1(km4)* mutants, SKN-1::GFP failed to accumulate in intestinal nuclei in response to IIS kinase RNAi (Figure 2B). Similarly, RNAi of the IIS kinases did not allow nuclear accumulation of a SKN-1 mutant in which two key p38 phosphorylation sites had been altered (Figure S3). We conclude that the SKN-1 nuclear accumulation that is allowed when IIS is reduced is appropriately dependent upon p38 MAPK signaling.

We next investigated whether SKN-1 is phosphorylated by the downstream IIS kinases AKT-1, -2, and SGK-1. Analysis using SCANSITE identified seven potential AKT phosphorylation motifs within SKN-1C (Table S1). Four additional potential AKT phosphorylation motifs were detected within the unique N-terminal 90 amino acids of SKN-1A (SKN-1A(1–90); Table S1), which is predicted by expressed sequence tag analysis to be the most abundant SKN-1 isoform (<http://www.wormbase.org>). AKT-1::GFP, AKT-2::GFP, and SGK-1::GFP that were purified from WT *C. elegans* (Hertweck et al., 2004) each phosphorylated full-length SKN-1C in vitro at least as robustly as they phosphorylated DAF-16 (Figure 2C). In contrast, SKN-1A(1–90) was phosphorylated by AKT-1 and AKT-2 but not SGK-1 (Figure 2D), consistent with the previous finding that AKT-1, -2, and SGK-1 differ with respect to their phosphorylation sites within the mammalian DAF-16 homolog FOXO3a (Brunet et al., 2001). Significantly, SKN-1 was not phosphorylated robustly by these same kinases if they were purified from *daf-2(e1370)* animals (Figures 2C and 2D). We conclude that SKN-1 is a direct substrate of the downstream IIS kinases, and that their phosphorylation of SKN-1 depends upon signaling through DAF-2.

We then performed a partial mapping of IIS kinase phosphorylation sites in SKN-1. Within SKN-1A(1–90), Ala substitution at the strongly predicted AKT site Ser12 reduced its phosphorylation by AKT-2 dramatically, (Figure 2D) identifying Ser12 as an AKT-2 site. AKT-1 phosphorylation was only slightly reduced by this mutation but was substantially decreased by concurrent substitution of Ser12 and each of three other sites within this region (Figure 2E). AKT and SGK also preferentially phosphorylated




Figure 2. DAF-2 Signaling Inhibits Nuclear Accumulation of SKN-1 Directly, and Independently of DAF-16

(A) *akt-1;akt-2* or *sgk-1* RNAi allows *daf-16*-independent accumulation of SKN-1 in intestinal nuclei. Expression of *SKN-1op::GFP* was monitored. *akt-1* or *akt-2* RNAi had a similar effect to combined *akt-1;akt-2* RNAi (data not shown).

(B) RNAi depletion of the downstream IIS kinases causes *sek-1*-dependent accumulation of SKN-1 in intestinal nuclei. Expression of *SKN-1B/C::GFP* was monitored.

(C) SKN-1C is phosphorylated by AKT-1, -2, and SGK-1 comparably to DAF-16, dependent upon *daf-2*. Here and in (D)–(G), AKT-1::GFP, AKT-2::GFP, and SGK-1::GFP were immunoprecipitated from WT or *daf-2(e1370)* *C. elegans* lysates with anti-GFP antibody, quantitated by western blotting with anti-GFP, and used to phosphorylate purified GST-fused SKN-1 or DAF-16 proteins with [γ - 32 P]ATP. GST controls are shown in Figure S4B. These kinases appropriately required activation in *C. elegans*, as they did not phosphorylate SKN-1 after they had been expressed in *E. coli* (Figure S4A).

(D) The unique 90 aa region at the SKN-A N terminus (SKN-1A(1–90)) is phosphorylated by the AKT kinases but not SGK-1, dependent upon *daf-2*.

(E) Partial mapping of AKT-1 sites within SKN-1A(1–90).

(F) AKT and SGK phosphorylate particular regions of SKN-1C. Here and in (G), residues are numbered as in SKN-1A, so that the first residue of SKN-1C is numbered as 91.

(G) Partial mapping of AKT-1, -2, and SGK-1 sites within SKN-1C fragments.

(H and I) Substitution of a strongly predicted AKT site within SKN-1A(1–90), Ser12 (Table S1), results in nuclear accumulation of SKN-1A::GFP. In some lines low levels of SKN-1::GFP were observed in additional head neurons, hypodermis, and other tissues, particularly if Ser12 was mutated (data not shown). p values were derived from a χ^2 test.

different regions or individual sites within the remainder of SKN-1, which corresponds to the SKN-1C isoform (Figures 2F and 2G; Table S1). Together, our data show that AKT-1, -2, and SGK-1 phosphorylate SKN-1 at multiple overlapping and distinct sites.

To investigate whether SKN-1 is regulated directly by IIS kinase phosphorylation in vivo, we substituted Ala at its most strongly predicted AKT site (Ser12 of SKN-1A) (Table S1), within the *SKN-1op::GFP* transgene. In each transgenic line examined, this substitution resulted in dramatic constitutive intestinal nuclear accumulation of SKN-1A (Figures 2H and 2I). This finding shows that SKN-1A is expressed in the intestine, and that the AKT phosphorylation site Ser12 is essential for its appropriate regulation. This strongly supports the model that signaling through the DAF-2 pathway directly inhibits SKN-1 in vivo.

Inhibition of SKN-1 Target Genes by IIS

Given that IIS directly inhibits the SKN-1 protein in the intestine, it follows that IIS should also reduce expression of SKN-1 target genes. To test this idea, we first asked how reductions in IIS affect expression of two SKN-1 target reporter transgenes (Figures 3A and 3B). In the intestine, SKN-1 directly induces expression of *gcs-1* (γ -glutamyl cysteine synthetase), which encodes an enzyme rate-limiting for glutathione biosynthesis (Figure 3A) (An and Blackwell, 2003). In the intestine a *gcs-1::GFP* transcriptional reporter is generally expressed at very low levels in the absence of stress but was upregulated by *akt-1;akt-2* or *sgk-1* RNAi (Figure 3C). This induction was abolished by mutation of an important SKN-1 binding site (*gcs-1 Δ 2mut3*; Figure 3A) but not by absence of *daf-16* (Figure 3C), indicating that SKN-1 but not DAF-16 was required for IIS kinase RNAi to activate the *gcs-1* promoter. Interestingly, *gcs-1* was not induced by *daf-2* RNAi, implying that its regulation by AKT and SGK-1 might involve an additional input (data not shown). In contrast, a translational fusion reporter for the SKN-1 target gene *gst-7* (glutathione S-transferase; Figure S5) was comparably induced in the intestine by either *daf-2* or IIS kinase RNAi (Figures 3B and 3D).

We next used quantitative (Q)-PCR to analyze how *daf-2* mutation affected the levels of mRNA expressed from a set of known or predicted SKN-1 target phase 2 genes (An and Blackwell, 2003; Kell et al., 2007). As would be expected, SKN-1 was not required for *daf-2* mutation to induce expression of a negative control for our experiments, the DAF-16 target *sod-3* (Figure 3E) (Antebi, 2007). The levels of *gcs-1* mRNA were largely unaffected in this *daf-2* mutant (Figure 3F), in agreement with our reporter data and the high levels of *skn-1*-independent *gcs-1* expression seen in the pharynx (An and Blackwell, 2003). In contrast, four different *gst* genes were strongly upregulated by *daf-2* mutation in a manner that was substantially or completely dependent upon *skn-1*. (Figures 3G–3J). DAF-16 contributed to induction of some of these genes, but the SKN-1 target *gst-10* was upregulated by *daf-2* mutation independently of *daf-16* (Figures 3G–3J). Together, our results indicate that when DAF-2 pathway signaling is reduced, SKN-1 induces a gene expression program that overlaps only partially with that of DAF-16.

skn-1 Contributes to the Increased Longevity and Stress Resistance Associated with Reduced IIS


Having established that DAF-2 and IIS inhibit SKN-1 directly, and thereby suppress SKN-1-dependent gene expression, we next

asked whether SKN-1 contributes to phenotypes that are associated with decreased IIS. We engineered *daf-2;skn-1* double mutants that each carried one of three *skn-1* mutations, along with one of two *daf-2* alleles. Each *skn-1* mutation (Figure 1B) results in reduced stress resistance and longevity (Figure 4; Table 1) (An and Blackwell, 2003), failure of DR to extend life span (Bishop and Guarente, 2007), and maternally derived embryonic lethality (Bowerman et al., 1992). The class 2 *daf-2* mutant *e1370* is stress resistant and lives much longer than WT at 20°C (Figures 4A and 4C), but this increased life span is accompanied by various dauer-like abnormalities (Gems et al., 1998; Kimura et al., 1997). The class 1 *daf-2* mutation *e1368* increases life span to a lesser extent (Figure 4D), but these animals appear essentially normal. We also tested how *skn-1* mutations affect similar phenotypes caused by RNAi of the IIS downstream kinases. At 25°C, *sgk-1* RNAi increases both oxidative stress resistance and life span, dependent upon *daf-16* (Hertweck et al., 2004). Previously, life span was not extended when *akt-1* RNAi was performed in the *akt-2(ok393)* mutant (Hertweck et al., 2004), but we have now observed that *akt-1;akt-2* RNAi significantly increases stress resistance and life span in an RNAi hypersensitive strain (Figures 4B and 4F; Table 1). This suggests that AKT-1, -2, and SGK-1 may contribute comparably to the effects of DAF-2 signaling on stress resistance and longevity.

Initially, we tested whether the IIS-associated phenotypes of dauer formation and elevated stress resistance require *skn-1*. *skn-1* RNAi did not prevent the constitutive dauer entry associated with the *daf-2(e1370)* mutation, or *akt-1;akt-2* RNAi (Supplemental Data). In contrast, and consistent with its role in phase 2 detoxification, mutations in *skn-1* dramatically suppressed the increases in oxidative stress resistance seen in each of these contexts of reduced IIS, and also with *sgk-1* RNAi (Figures 4A and 4B).

We then investigated how *skn-1* contributes to the increased longevity deriving from decreased *daf-2* function. Only *skn-1(zu129)* suppressed the increased longevity of the class 2 *daf-2* allele *e1370* (Figure 4C; Table 1). In contrast, all three *skn-1* mutations significantly reduced the life span of the class 1 *daf-2* mutant *e1368* (Figure 4D; Table 1). These *skn-1* mutations also altered the shape of the *daf-2(e1368)* survival plot, so that the longest-living double mutant animals appeared to be less dependent upon *skn-1* than the remaining population (Figure 4D). Accordingly, when we excluded the 20% most long-lived animals from statistical analysis, two of three *skn-1* mutations affected death rate more severely in *daf-2(e1368)* animals than in WT (Figure S7; Tables S3 and S4). Importantly, an RNAi-sensitive genetic background *skn-1* RNAi decreased the life span of *daf-2(e1368)* but not control animals (Figure 4E; Tables 1 and S4), indicating that this *daf-2* mutant is more susceptible than WT to a reduction in SKN-1 levels. *daf-2(e1368)* life span was similarly decreased by either *skn-1* or *skn-1a/c* RNAi (Figure 4E; Tables 1 and S4), indicating that the SKN-1 A and/or C isoforms are required for the longevity associated with *daf-2(e1368)*. *daf-2(e1368)* life span was also decreased by feeding of *skn-1* RNAi from the beginning of adulthood (Figure S8), indicating an effect on aging, not development.

Mutations in *skn-1* also suppressed the life-span extensions associated with reductions in IIS kinase levels. Each of two *skn-1* alleles that we analyzed (*zu67* and *zu129*) completely eliminated



the dramatic increase in life span deriving from *sgk-1* RNAi (Figure 4G; Tables 1 and S4). In addition, two of three *skn-1* alleles significantly reduced the life spans of *akt-1;akt-2* RNAi animals (Figure 4F; Table 1). We conclude that SKN-1 makes important contributions to the increases in stress resistance and longevity that are associated with reductions in IIS activity.

Next, we investigated whether *skn-1* affects the long life spans of animals with reduced mitochondrial electron transport chain activity to find out whether *skn-1* mutations might cause a general sickness that limits life span (Dillin et al., 2002; Lee et al., 2003; Rea et al., 2007). In agreement with a recent report (Rea et al., 2007), a *skn-1* mutation did not reduce the large increases in life span associated with RNAi against either cytochrome *c* reductase (*cyc-1*) or cytochrome *c* oxidase (*cco-1*) (Figures 4H and S9; Table 1). This indicates that while *skn-1* is important for life span in long-lived animals in which IIS is reduced, lack of *skn-1* does not limit longevity nonspecifically.

SKN-1 Promotes Longevity

The involvement of *skn-1* in the longevity effects of IIS led us to investigate whether SKN-1 has an independent pro-longevity function. Previously, life span was modestly increased by transgenic overexpression of either WT or constitutively nuclear forms of DAF-16 (Henderson and Johnson, 2001; Lin et al., 2001). To investigate whether modulation of SKN-1 expression can extend life span, we first analyzed otherwise WT strains in which either the *SKN-1B/C::GFP* or *SKN-1B/C S393A::GFP* transgenes were present. The *SKN-1B/C S393A::GFP* transgene, which expresses constitutively nuclear SKN-1 in the intestine, increased mean life span variably but consistently (by 5%–21%; Figure 5A; Table 2). A modest increase was also seen with *SKN-1B/C::GFP* (*Ex001*; Figure 5B; Table 2).

We had previously observed that *skn-1* transgenes not only rescue the oxidative stress sensitivity of *skn-1(zu67)* mutants but also increase their stress resistance to well beyond that of WT (An et al., 2005). Similarly, strains in which *skn-1(zu67)* was rescued with these transgenes also lived longer than wild-type animals that carried the same transgene arrays (arrays *Ex001* and *Ex008*, Figures 5A and 5B; Table 2). This difference was corroborated in independent strains that were generated by breeding the *Ex001[SKN-1B/C::GFP]* array from the WT into *skn-1(zu67)* and *skn-1(zu129)* (Figures S10A, S10B, and S11), and in an additional *skn-1(zu67)* strain carrying *SKN-1B/C S393A::GFP* (*Ex020*, Figure 5B; Table 2). Perhaps an optimal window of SKN-1 expression is important for longevity because we found that expression of SKN-1 from high-copy arrays can be toxic (data not shown). Significantly, rescued *skn-1(zu67)* *Ex001[SKN-1B/C::GFP]* animals maintained the ability to move longer than WT (Figure S10C), indicating that transgenic *skn-1* expression delayed the aging pro-

cess. RNAi that was targeted to transgenic SKN-1C (*skn-1a/c* RNAi, Figure 1B) eliminated expression of intestinal nuclear SKN-1 from *SKN-1B/C S393A::GFP* (Figure S12) and suppressed the extended life span of a transgenically rescued *skn-1(zu67)* strain (Figure 5C). Taken together, our results indicate that SKN-1 promotes longevity, an effect that appears to require the SKN-1C isoform.

We next investigated whether *daf-16* is required for transgenic SKN-1 to increase life span because of the general importance of *daf-16* for *C. elegans* longevity and because SKN-1 and DAF-16 overlap in regulating some target genes (Figures 3E–3J). When we bred a SKN-1 S393A transgenic array (*Ex008*) into the null mutant *daf-16(mgDf47)*, life span was modestly but significantly increased in two of three experiments (Figure 5D; Tables 2 and S6). Similarly, after *daf-16* RNAi the life spans of *skn-1(zu67)* animals were reduced, but presence of an array that expresses constitutively nuclear SKN-1(S393A) (*Ex008* or *020*) still increased *skn-1(zu67)* life span to beyond that of WT (Figure 5E; Table 2). Interestingly, *daf-16* RNAi completely suppressed the life-span extension associated with transgenic WT SKN-1 (Figure S13; Table 2), suggesting that reductions in DAF-16 levels not only reduce life span but also might free up negative regulatory mechanisms that act on SKN-1. Taken together, our data suggest that SKN-1 can promote longevity independently of DAF-16, provided that SKN-1 is released from negative regulation in the intestine.

DISCUSSION

It is well established that FOXO proteins are inhibited directly by the IIS pathway, and that the FOXO protein DAF-16 is required for the increased longevity and stress resistance that result from decreased IIS in *C. elegans* (Kenyon, 2005). This has made it difficult to imagine that another transcription factor might be regulated similarly by IIS in parallel to FOXO and could play a major role in IIS-regulated processes. However, here we have identified SKN-1 as the second transcription factor that is known to be inhibited directly by IIS. We have also shown that IIS inhibits SKN-1 target genes and that SKN-1 contributes to the stress resistance and longevity phenotypes of reduced IIS and functions as a pro-longevity factor.

Direct Regulation of SKN-1 by IIS

Multiple lines of evidence indicate that IIS inhibits SKN-1 in the intestine directly (Figure 1A). Reductions in IIS activity resulted in SKN-1A and SKN-1C being present in intestinal nuclei in the absence of either stress or DAF-16 and in induction of SKN-1 target genes (Figures 1E, 2A, 2B, and 3). SKN-1 isoforms were phosphorylated by AKT-1, -2, and SGK-1 at multiple sites, and

(C) *gcs-1::GFP* is induced in the intestine independently of *daf-16* in response to *akt-1;akt-2* and *sgk-1* RNAi. Here and in (D), results of multiple experiments performed on L4 larvae have been combined and scored as described in the Supplemental Experimental Procedures and An and Blackwell (2003) and An et al. (2005). p values were derived from a χ^2 test. *akt-1* and *akt-2* RNAi were also performed separately and shown to have a similar effect to combined *akt-1;akt-2* RNAi (data not shown).

(D) *GST-7::GFP* is induced in response to *daf-2*, *akt-1;akt-2*, or *sgk-1* RNAi.

(E–J) SKN-1- and DAF-16-responsive genes that are upregulated in *daf-2(e1370)* L4 larvae were identified by quantitative PCR, using actin (*act-1*) as a control. The level of *skn-1* mRNA in these samples was also measured (Figure S6). One representative experiment is shown, error bars represent the SEM, and p values were derived from a Student's t test.




Figure 4. *skn-1* Mutations Suppress the Longevity and Stress Resistance Phenotypes Conferred by Reduced IIS

(A) *skn-1* contributes to the oxidative stress resistance phenotype of *daf-2(e1370)* mutants. A representative experiment is shown in which animals were exposed to 7.5 mM *t*-butyl hydrogen peroxide. Here and in (B), error bars represent the SEM.

(B) *skn-1* contributes to the stress resistance phenotypes conferred by *sgk-1* or *akt-1;akt-2* RNAi. Animals were scored after 5 days exposure to 150 mM paraquat. Data were combined from at least three experiments with $n > 90$ for each group.

(C) Life-span analysis of *daf-2(e1370); skn-1* mutants.

mutation of an AKT site (Ser12) led to SKN-1A being present in intestinal nuclei constitutively (Figures 2H and 2I). The AKT and SGK kinases differed in their preferences for sites within SKN-1, consistent with previous analyses of their phosphorylation of FOXO (Brunet et al., 2001). The complexity of this regulation could potentially allow SKN-1 to integrate inputs from multiple signals or molecular interactions.

We also observed that SKN-1 contributes significantly to some IIS-associated phenotypes. *skn-1* mutations dramatically suppressed the increases in stress resistance deriving from *sgk-1* RNAi, *akt-1;akt-2* RNAi, and the *daf-2(e1370)* mutation (Figures 4A and 4B). Lack of *skn-1* completely suppressed the increased life span deriving from *sgk-1* RNAi, reduced the extended longevity of *akt-1;akt-2* RNAi animals (Figures 4F and 4G), and suppressed the increased life span of the class 1 *daf-2* allele *e1368* (Figure 4D). Importantly, *skn-1* RNAi reduced the life span of *daf-2(e1368)* but not WT animals (Figure 4E), indicating that *daf-2(e1368)* was more severely affected than WT by partial loss of *skn-1* function. In contrast, *skn-1* mutation did not suppress the long life associated with RNAi of mitochondrial respiratory chain components (Figures 4H and S9), indicating that lack of *skn-1* does not reduce life span nonspecifically. *skn-1* mutations also did not consistently suppress the life-span extension of *daf-2(e1370)* mutants (Figure 4C). However, dauer-like phenotypes are associated with this class 2 allele (Gems et al., 1998), suggesting that this difference might reflect the apparent dispensability of SKN-1 for dauer formation (Supplemental Data).

The simplest model to explain the contributions of SKN-1 to IIS-associated longevity is that they derive from phase 2 detoxification gene expression in the intestine. Our finding that *skn-1a/c* RNAi suppresses *daf-2(e1368)* longevity (Figure 4E) is consistent with this idea, although we cannot exclude that other tissues or SKN-1 functions are involved. We conclude that *daf-16* is necessary but not sufficient to confer the increases in stress resistance and longevity that are associated with reduced IIS, and that these increases also involve *skn-1*.

In contrast to the regulation of SKN-1 by IIS in the intestine, *daf-2* mutation did not obviously affect SKN-1 levels in the ASI neurons (Figure 1C; not shown), in which SKN-1B is required for DR to extend life span (Bishop and Guarente, 2007). However, we cannot exclude that IIS might subtly influence the levels or activity of endogenous SKN-1 in ASI. This possibility could help resolve the apparent paradox that in *C. elegans* most DR protocols extend life span independently of *daf-16*, and therefore seemingly without input from IIS (Antebi, 2007; Bishop and Guarente, 2007; Greer et al., 2007; Houthoofd et al., 2003; Kenyon, 2005; Panowski et al., 2007).

SKN-1 is the sequence and functional counterpart to the mammalian NF-E2-related factor (Nrf) proteins (An and

Blackwell, 2003), which orchestrate the phase 2 detoxification response (Hayes and McMahon, 2001). Our results predict that the conserved IIS pathway might also regulate mammalian Nrf proteins, which include potential AKT sites (Table S1). Why would it be advantageous for IIS to decrease stress resistance? Many signaling cascades are influenced by redox status (Tonks, 2005). In particular, IIS is attenuated by protein tyrosine phosphatase (PTP)1B, which is inhibited by reactive oxygen species (ROS) via the oxidation of conserved cysteine residues (Goldstein et al., 2005; Tonks, 2005). Thus, by inhibiting DAF-16/FOXO and SKN-1/Nrf, IIS could suppress antioxidant defenses that would lead to a reducing environment, and that in turn might decrease IIS effectiveness. This is consistent with evidence that insulin induces starved cells to produce a “burst” of ROS, and that IIS functions more effectively under oxidative conditions (Goldstein et al., 2005).

Pro-longevity Functions of SKN-1

Our finding that transgenic SKN-1 expression extends life span (Figures 5A and 5B) demonstrates that *skn-1* has an independent pro-longevity function. This activity appears to require the action of the intestinal SKN-1C isoform because it is suppressed by *skn-1a/c* RNAi (Figure 5C). This pro-longevity activity of SKN-1 thus appears to be distinct from its ASI-mediated role in DR-induced longevity (Bishop and Guarente, 2007). The observation that transgenic SKN-1 increased life span more robustly in *skn-1* mutants than in the WT (Figures 5A and 5B) suggests that the systemic effects of SKN-1 expression may be complex. However, taken together, our data demonstrate that SKN-1 levels can be adjusted to extend life span and delay aging. We propose that the ancestral function of SKN-1 in phase 2 detoxification can be modulated by either reductions in IIS or adjustment of SKN-1 expression to extend life span, raising the exciting possibility that this detoxification system might be of general and conserved importance for longevity.

In the context of reduced IIS, SKN-1 and DAF-16 appeared to have both distinct and overlapping functions (Figure 3). *daf-16* contributed to expression of some *skn-1*-dependent phase 2 detoxification genes (Figures 3E–3I), but other SKN-1 targets were *daf-16* independent (Figure 3J), consistent with our finding that transgenic expression of SKN-1 S393A increased longevity independently of *daf-16* (Figures 5D and 5E). Our data indicate that SKN-1 has important activities that are largely independent of DAF-16, making it of intense interest to dissect the extent to which these two transcription factors cooperate or act independently under normal, stress, and reduced IIS conditions. It is interesting that transgenic overexpression of neither DAF-16 or SKN-1 recapitulates the dramatic increases in longevity seen with reductions in DAF-2 function (Lin et al., 2001; Figure 5). One possibility is that

(D) *skn-1* suppresses the longevity of *daf-2(e1368)* mutants.

(E) In the RNAi-sensitive *rrf-3* background (Simmer et al., 2002), the life-span increase seen in *daf-2(e1368)* is comparably suppressed by either *skn-1* or *skn-1a/c* RNAi. RNAi bacteria were provided from the L1 stage.

(F and G) *skn-1* mutations suppress the life-span extension conferred by *akt-1;akt-2* or *sgk-1* RNAi. Survival assays were carried out at 25°C, are from adulthood, and represent the combined data from at least two experiments.

(H) *skn-1* mutation does not suppress the life-span extension conferred by *cyc-1* RNAi. The life-span experiments in (C) and (D) were carried out on OP50, but all others were performed on HT115 lawns. Life-span data in this Figure are summarized in Table 1. Unless otherwise indicated, the analyses shown were performed at 20°C and measured from hatching.

Table 1. *skn-1* Contributes to the Life-Span Extension Resulting from Reduced IIS

| Strain | Mean Life | Median | 75 th | 25 th | p Value | n | No. of Exp. |
|--|-------------|-----------|------------------|------------------|---------------------|-------|-------------|
| | Span ± SEM | | Percentile | Percentile | | | |
| | 20°C (days) | Life Span | 20°C (days) | 20°C (days) | Control | | |
| N2 | 20.44 ± 0.3 | 21 | 23 | 19 | — | 90/90 | 1 |
| <i>daf-2(e1368)</i> | 29.42 ± 0.3 | 31 | 31 | 29 | <.0001 ^a | 91/91 | 1 |
| <i>daf-2(e1370)</i> | 39.39 ± 0.4 | 39 | 43 | 37 | <.0001 ^a | 88/88 | 1 |
| <i>daf-16(mgDf47)</i> | 16.80 ± 0.3 | 17 | 19 | 15 | <.0001 ^a | 50/50 | 1 |
| <i>daf-16(mgDf47); daf-2(e1370)</i> | 20.01 ± 0.1 | 21 | 21 | 19 | <.0001 ^b | 77/80 | 1 |
| <i>skn-1(zu67)</i> | 15.60 ± 0.3 | 17 | 17 | 15 | <.0001 ^a | 73/73 | 1 |
| <i>skn-1(zu129)</i> | 17.67 ± 0.2 | 17 | 19 | 17 | <.0001 ^a | 90/90 | 1 |
| <i>skn-1(zu135)</i> | 17.70 ± 0.3 | 17 | 19 | 17 | <.0001 ^a | 57/57 | 1 |
| <i>daf-2(e1368); skn-1(zu67)</i> | 22.63 ± 0.5 | 23 | 25 | 21 | <.0001 ^c | 60/60 | 1 |
| <i>daf-2(e1368); skn-1(zu129)</i> | 22.25 ± 1.1 | 19 | 29 | 15 | <.0001 ^c | 58/60 | 1 |
| <i>daf-2(e1368); skn-1(zu135)</i> | 19.83 ± 0.6 | 19 | 23 | 17 | <.0001 ^c | 58/59 | 1 |
| <i>daf-2(e1370); skn-1(zu67)</i> | 38.60 ± 1.0 | 41 | 43 | 35 | .0740 ^b | 60/60 | 1 |
| <i>daf-2(e1370); skn-1(zu129)</i> | 27.64 ± 0.7 | 27 | 31 | 23 | <.0001 ^b | 62/62 | 1 |
| <i>daf-2(e1370); skn-1(zu135)</i> | 33.16 ± 0.6 | 39 | 43 | 33 | .6016 ^b | 25/25 | 1 |
| <i>rrf-3(pk1426); control(RNAi)</i> | 19.53 ± 0.5 | 19 | 24 | 17 | — | 57/60 | 1 |
| <i>rrf-3(pk1426); skn-1(RNAi)</i> | 20.43 ± 0.6 | 21 | 24 | 15 | .1531 ^g | 60/60 | 1 |
| <i>rrf-3(pk1426); skn-1a/c(RNAi)</i> | 20.27 ± 0.6 | 21 | 24 | 15 | .2313 ^g | 59/60 | 1 |
| <i>rrf-3(pk1426); daf-2(e1368) control(RNAi)</i> | 38.62 ± 1.3 | 40 | 44 | 34 | <.0001 ^g | 52/55 | 1 |
| <i>rrf-3(pk1426); daf-2(e1368) skn-1(RNAi)</i> | 25.83 ± 1.3 | 21 | 36 | 17 | <.0001 ^d | 56/56 | 1 |
| <i>rrf-3(pk1426); daf-2(e1368) skn-1a/c (RNAi)</i> | 26.32 ± 1.3 | 24 | 36 | 19 | <.0001 ^d | 57/58 | 1 |
| N2 [rol-6]; control(RNAi) | 20.1 ± 0.3 | 20 | 22 | 18 | — | 84/84 | 1 |
| <i>skn-1(zu67); control(RNAi)</i> | 14.6 ± 0.4 | 14 | 16 | 12 | <.0001 ^e | 87/90 | 1 |
| N2 [rol-6]; <i>cyc-1(RNAi)</i> | 42.1 ± 0.9 | 42 | 45 | 40 | <.0001 ^e | 32/34 | 1 |
| <i>skn-1(zu67); cyc-1(RNAi)</i> | 41.9 ± 1.2 | 45 | 47 | 38 | .5594 ^f | 38/38 | 1 |
| N2 [rol-6]; <i>cco-1(RNAi)</i> | 31.52 ± 0.5 | 32 | 34 | 30 | <.0001 ^e | 67/69 | 1 |
| <i>skn-1(zu67); cco-1(RNAi)</i> | 35.13 ± 0.5 | 36 | 38 | 34 | <.0001 ^f | 46/46 | 1 |

| Strain | Mean Adult | Median | 75 th | 25 th | p Value | n | No. of Exp. |
|---|-----------------|-----------|------------------|------------------|---------------------|---------|-------------|
| | Life Span ± SEM | | Percentile | Percentile | | | |
| | 25°C (days) | Life Span | 25°C (days) | 25°C (days) | Control | | |
| <i>rrf-3(pk1426); control(RNAi)</i> | 12.90 ± 0.4 | 13 | 15 | 11 | — | 99/100 | 2 |
| <i>daf-16(mgDf47)</i> | 11.04 ± 0.2 | 11 | 13 | 9 | <.0001 ^g | 138/138 | 3 |
| <i>rrf-3(pk1426); akt1; akt-2(RNAi)</i> | 22.50 ± 0.7 | 22 | 28 | 17 | <.0001 ^g | 120/138 | 2 |
| <i>daf-16(mgDf47); rrf-3(pk1426); akt1; akt-2(RNAi)</i> | 11.03 ± 0.3 | 11 | 13 | 10 | <.0001 ^h | 123/123 | 3 |
| <i>rrf-3(pk1426); skn-1(zu67); control(RNAi)</i> | 7.20 ± 0.2 | 7 | 8 | 6 | <.0001 ^g | 106/106 | 2 |
| <i>rrf-3(pk1426); skn-1(zu129); control(RNAi)</i> | 9.30 ± 0.4 | 8 | 12 | 7 | <.0001 ^g | 100/100 | 2 |
| <i>rrf-3(pk1426); skn-1(zu135); control(RNAi)</i> | 12.00 ± 0.3 | 12 | 15 | 9 | 0.1012 ^g | 150/150 | 3 |
| <i>rrf-3(pk1426); skn-1(zu67); akt-1; akt-2(RNAi)</i> | 11.60 ± 0.6 | 11 | 17 | 6 | <.0001 ^h | 104/104 | 2 |
| <i>rrf-3(pk1426); skn-1(zu129); akt-1; akt-2(RNAi)</i> | 20.40 ± 1.0 | 21 | 28 | 12 | 0.9400 ^h | 101/104 | 2 |
| <i>rrf-3(pk1426); skn-1(zu135); akt-1; akt-2(RNAi)</i> | 17.69 ± 0.6 | 17 | 22 | 13 | <.0001 ^h | 110/110 | 3 |
| N2; control(RNAi) | 13.8 ± 0.3 | 14 | 16 | 12 | — | 110/114 | 2 |
| N2; <i>sgk-1(RNAi)</i> | 22.6 ± 0.4 | 23 | 25 | 21 | <.0001 ^e | 110/395 | 2 |
| <i>skn-1(zu67); sgk-1(RNAi)</i> | 8.2 ± 0.2 | 8 | 10 | 7 | <.0001 ⁱ | 128/128 | 2 |
| <i>skn-1(zu129); sgk-1(RNAi)</i> | 9.1 ± 0.2 | 8 | 11 | 7 | <.0001 ⁱ | 185/186 | 2 |

Corresponds to the data in Figures 4C–4H. SEM: standard error of the mean. 75th and 25th percentiles refer to the day at which 75% or 25% of the population is dead. n represents total number of animals dying of old age versus those in total experiment (combined between trials where appropriate). In the case of *sgk-1(RNAi)* the majority of exclusions are due to *Egl/bag* of worms phenotypes. p values were calculated as follows: ^aN2, ^b*daf-2(e1370)*, ^c*daf-2(e1368)*, ^d*rrf-3(pk1426); daf-2(e1368) control(RNAi)*, ^eN2 control(RNAi), ^fN2 with corresponding RNAi, ^g*rrf-3(pk1426) control(RNAi)*, ^h*rrf-3(pk1426) akt-1; akt-2(RNAi)*, ⁱ*sgk-1(RNAi)*. Individual experiments, an additional analysis of *daf-2(e1368)* and *daf-2(e1370)*, and other supporting data are in Table S5.




Figure 5. Transgenic Expression of SKN-1 Extends Life Span

(A) Life-span extension in otherwise WT animals carrying the *SKN-1B/C S393A::GFP* transgene, which expresses constitutively nuclear SKN-1C in the intestine (see text). These independent extrachromosomal arrays were created by injecting 2.5 ng/ μ l transgene DNA.

(B) Life-span extension in *skn-1(zu67)* animals carrying *skn-1* transgenes. The Ex001[*SKN-1B/C::GFP*], Ex008[*SKN-1B/C S393A::GFP*], and Ex020[*SKN-1B/C S393A::GFP*] extrachromosomal arrays were originally created in WT animals by injection of 2.5, 2.5, and 10 ng/ μ l transgene DNA, respectively, and then bred into *skn-1(zu67)*. The presence of the transgene did not dramatically affect the developmental timing, brood size, or age-specific fecundity of either the WT or *skn-1* mutants (Figure S14). The life spans of WT animals carrying Ex001[*SKN-1B/C::GFP*] is shown for comparison.

(C) SKN-1C contributes to the life span of *skn-1(zu67)* Ex001[*SKN-1B/C::GFP*] animals. Life span is reduced comparably by *skn-1* and *skn-1a/c* RNAi (Figure 1B).

(D) Presence of Ex008[*SKN-1B/C S393A::GFP*] increases the life span of *daf-16(mgDf47)* mutants.

(E) Transgene arrays that express constitutively nuclear SKN-1 (Ex008 or Ex020) increase *skn-1(zu67)* life span beyond WT when *daf-16* is depleted by RNAi. Life-span analyses were scored from hatching and performed at 20°C unless otherwise stated and are summarized in Table 2. The experiments in (A) and (B) were carried out on OP50, and the others on HT115. (C) and (D) are representative experiments, but otherwise data were combined from at least two experiments, within which all sample sets were analyzed in parallel.

(F) Crosstalk among *C. elegans* transcriptional regulators that increase longevity when expressed transgenically. DAF-16 physically interacts with SIR-2.1 through 14-3-3 proteins and is required for SIR-2.1 to extend life span (Berdichevsky et al., 2006; Tissenbaum and Guarente, 2001). HSF-1 and DAF-16 are predicted to bind to some of the same promoters and regulate some pro-longevity target genes together (Hsu et al., 2003). PHA-4 shares a common cooperating cofactor with DAF-16 (SMK-1), extends life span when DAF-16 is absent, and is required for DR to extend life span (Panowski et al., 2007). SKN-1 and DAF-16 are inhibited directly and in parallel by IIS (this work).

Table 2. Transgenic Expression of SKN-1 Extends Life Span

| Strain | Mean Life Span \pm SEM 20°C (days) | 75 th Percentile 20°C (days) | p value against control | n | No. of Exp. |
|--|---|---|-------------------------------|---------|----------------|
| <i>N2 Ex[rol-6]</i> | 19.57 \pm 0.2 | 21 | — | 129/130 | 2 |
| <i>N2 Ex008[SKN-1 S393A::GFP] 2.5 ng/μl</i> | 23.62 \pm 0.3 | 26 | <.0001 ^a | 78/78 | 2 |
| <i>N2 Ex021[SKN-1 S393A::GFP] 2.5 ng/μl</i> | 21.80 \pm 0.2 | 23 | <.0001 ^a | 117/122 | 2 |
| <i>N2 Ex022[SKN-1 S393A::GFP] 2.5 ng/μl</i> | 21.10 \pm 0.3 | 23 | <.0001 ^a | 62/62 | 2 |
| <i>N2 Ex023[SKN-1 S393A::GFP] 2.5 ng/μl</i> | 20.89 \pm 0.3 | 23 | <.0001 ^a | 96/96 | 2 |
| <i>N2 Ex024[SKN-1 S393A::GFP] 2.5 ng/μl</i> | 20.66 \pm 0.4 | 23 | <.0001 ^a | 63/63 | 2 |
| <i>N2 Ex[rol-6]</i> | 19.52 \pm 0.2 | 21 | — | 236/243 | 3 |
| <i>skn-1(zu67) Ex[rol-6]</i> | 15.51 \pm 0.3 | 18 | <0.0001 ^a | 96/96 | 2 |
| <i>N2 Ex001[SKN-1B/C::GFP] 2.5 ng/μl</i> | 22.14 \pm 0.3 | 25 | <.0001 ^{b,c} | 146/148 | 2 |
| <i>skn-1(zu67) Ex001[SKN-1B/C::GFP] 2.5 ng/μl</i> | 23.53 \pm 0.6 | 26 | <.0001 ^{b,c} | 48/51 | 2 |
| <i>skn-1(zu67) Ex008[SKN-1B/C S393A::GFP] 2.5 ng/μl</i> | 25.74 \pm 0.5 | 29 | <.0001 ^{b,c} | 67/70 | 2 |
| <i>skn-1(zu67) Ex020[SKN-1B/C S393A::GFP] 10 ng/μl</i> | 27.16 \pm 0.6 | 29 | <.0001 ^{b,c} | 30/40 | 1 |
| <i>N2 Ex[rol-6]; control(RNAi)</i> | 20.60 \pm 0.5 | 23 | — | 50/50 | 1 |
| <i>N2 Ex[rol-6]; skn-1(RNAi)</i> | 19.91 \pm 0.4 | 21 | 0.0699 ^f | 60/60 | 1 |
| <i>N2 Ex[rol-6]; skn-1a/c(RNAi)</i> | 20.37 \pm 0.4 | 23 | 0.7880 ^f | 59/60 | 1 |
| <i>skn-1(zu67) Ex001[SKN-1B/C::GFP] 2.5 ng/μl; control(RNAi)</i> | 24.20 \pm 0.4 | 27 | <.0001 ^f | 62/64 | 1 |
| <i>skn-1(zu67) Ex001[SKN-1B/C::GFP] 2.5 ng/μl; skn-1(RNAi)</i> | 21.58 \pm 0.4 | 23 | <.0001 ^d | 84/84 | 1 |
| <i>skn-1(zu67) Ex001[SKN-1B/C::GFP] 2.5 ng/μl; skn-1a/c(RNAi)</i> | 21.06 \pm 0.5 | 25 | 0.0005 ^d | 69/69 | 1 |
| <i>N2 Ex[rol-6]</i> | 20.60 \pm 0.5 | 23 | — | 50/50 | 1 |
| <i>daf-16(mgDf47)</i> | 15.87 \pm 0.3 | 17 | <.0001 ^f | 8/61 | 1 |
| <i>daf-16(mgDf47) Ex008[SKN-1 S393A::GFP] 2.5 ng/μl</i> | 18.88 \pm 0.3 | 21 | <.0001 ^e | 48/48 | 1 |
| <i>N2 Ex[rol-6]; control(RNAi)</i> | 20.02 \pm 0.2 | 22 | — | 256/259 | 4 |
| <i>N2 Ex[rol-6]; daf-16(RNAi)</i> | 17.77 \pm 0.1 | 19 | <.0001 ^f | 266/266 | 4 |
| <i>skn-1(zu67) Ex[rol-6]; control(RNAi)</i> | 17.38 \pm 0.3 | 20 | <.0001 ^f | 145/146 | 3 |
| <i>skn-1(zu67) Ex[rol-6]; daf-16(RNAi)</i> | 16.59 \pm 0.3 | 19 | 0.0240 ^g | 142/145 | 3 |
| <i>skn-1(zu67) Ex001[SKN-1b/c::GFP] 2.5 ng/μl; control(RNAi)</i> | 22.95 \pm 0.5 | 26 | <.0001 ^g | 62/62 | 2 |
| <i>skn-1(zu67) Ex001[SKN-1b/c::GFP] 2.5 ng/μl; daf-16(RNAi)</i> | 16.53 \pm 0.3 | 18 | 0.1655 ^h | 59/59 | 2 |
| <i>skn-1(zu67) Ex008[SKN-1 S393A::GFP] 2.5 ng/μl; control(RNAi)</i> | 24.28 \pm 0.6 | 28 | <.0001 ^g | 64/76 | 3 |
| <i>skn-1(zu67) Ex008[SKN-1 S393A::GFP] 2.5 ng/μl; daf-16(RNAi)</i> | 21.12 \pm 0.4 | 23 | <.0001 ^{h,i} | 76/87 | 3 |
| <i>skn-1(zu67) Ex020[SKN-1 S393A::GFP] 10 ng/μl; control(RNAi)</i> | 25.10 \pm 0.6 | 28 | <.0001 ^g | 121/132 | 3 |
| <i>skn-1(zu67) Ex020[SKN-1 S393A::GFP] 10 ng/μl; daf-16(RNAi)</i> | 21.39 \pm 0.4 | 24 | <.0001 ^{h,i} | 124/140 | 3 |

Corresponds to the data in Figures 5A–5E. p values were calculated as follows: ^a*N2 Ex[rol-6]*, ^b*skn-1(zu67)[rol-6]*, ^csignificantly enhanced compared to *N2 [rol-6]* ($p < 0.0001$ in each case), ^d*skn-1(zu67) Ex001 control(RNAi)*, ^e*daf-16(mgDf47) control(RNAi)*, ^f*N2 Ex[rol-6] control(RNAi)*, ^g*skn-1(zu67)[rol-6] control(RNAi)*, ^h*skn-1(zu67)[rol-6] daf-16(RNAi)*, ⁱcompared to *N2 Ex[rol-6] control(RNAi)* ($p = 0.0021$ for *skn-1(zu67) Ex008* and $p = 0.0001$ for *skn-1(zu67) Ex020*). Individual experiments and additional analysis of *daf-16 Ex008[SKN-1S393A::GFP]* are in Table S6. Tabulated as in Table 1.

simultaneous overexpression of both transcription factors is necessary, although modifications of these proteins or other signals associated with decreased IIS might also be required. Our observation that *daf-16* RNAi suppressed life-span extension by the WT *SKN-1B/C::GFP* transgene (Figure S13) suggests that DAF-16 may influence processes that regulate SKN-1. This could involve DAF-16 regulating genes that affect SKN-1, but a simpler model is that SKN-1 and DAF-16 compete for binding to negative regulators that include AKT-1, -2, and SGK-1 (Figure 1A). Thus, decreasing DAF-16 levels would increase the ratio of negative regulators available to sequester SKN-1 in the cytoplasm.

Our results add SKN-1 to the small group of *C. elegans* transcription regulators that extend longevity when expressed trans-

genically (Figure 5F). SIR-2.1, PHA-4, DAF-16, and SKN-1 or their orthologs have all been shown to influence aspects of metabolism and thereby presumably may affect the rates of damage that lead to aging (Accili and Arden, 2004; Bishop and Guarente, 2007; Kenyon, 2005; Panowski et al., 2007; van der Horst and Burgering, 2007). Importantly, SKN-1, HSF-1, and each of these other proteins have also been implicated in resistance to free radical or other stresses (Figure 5F; see text), supporting the idea that stress defenses and detoxification are fundamentally important for longevity. A remarkable web of functional interactions among these factors has been uncovered, including our demonstration here that DAF-16 and SKN-1 are regulated in parallel by IIS (Figure 5F). These interactions could

be important for coordinating how these proteins respond in different tissues to various scenarios of nutrient availability and stress. As these regulators are also evolutionarily conserved, it seems likely that a deeper understanding of interactions among them will be broadly relevant to elucidating how these proteins influence metabolism, stress resistance, and possibly longevity across species.

EXPERIMENTAL PROCEDURES

All **Supplemental Data** and experimental procedures, including details of cloning, strains, RNAi, GFP scoring system, and assays for life span, movement, and reproduction are available online.

In Vitro Kinase Assay

Kinases were isolated from sonicated worm lysates of *N2 Ex[AKT-1::GFP]*, *N2 Ex[AKT-2::GFP]*, and *N2 Ex[AKT-1::GFP]* (Hertweck et al., 2004), using a GFP antibody bound to protein A/G agarose beads. These purified kinases were used to phosphorylate 6 μ g of bacterially expressed GST-SKN-1 fusion proteins in the presence of 6 μ Ci [γ P³²]ATP. After phosphorylation, samples were washed in 50 mM Tris/HCl (pH 8.0), 100 mM NaCl, 10% glycerol, and 1% Triton X-100 and analyzed by SDS-PAGE and autoradiography.

RNAi

Feeding RNAi was carried out as previously described, with empty pL4440 as the control (**Supplemental Experimental Procedures**). To avoid maternal lethality in experiments using *skn-1* or *skn-1a/c* RNAi, animals were hatched on HT115 bacteria expressing control RNAi and were transferred to *skn-1* RNAi at the L1 stage or on the first day of adulthood. For life-span experiments that required feeding of other RNAi, animals were exposed to bacteria expressing the appropriate construct continually starting from hatching.

Life-Span Analysis

Prior to experiments, all animals were maintained at the permissive temperature and grown for at least two generations in the presence of food to assure health. Life-span assays were performed essentially as described (Hsin and Kenyon, 1999; **Supplemental Experimental Procedures**). Survival plots, p values (Log-Rank), and proportional hazards were determined using JMP software, version 5.1.

Oxidative Stress Resistance Assay

To assess stress resistance, young adults were transferred to plates that contained 7.5 mM *t*-butyl hydrogen peroxide (Sigma) or 150 mM paraquat (Sigma) in nematode growth media. These plates had been seeded with OP50 or HT115 expressing the appropriate dsRNAs. Animals were incubated on these plates at 20°C and periodically scored for survival.

RNA Isolation and Quantitative PCR

Animals were picked to clean plates to minimize contamination, then total RNA was extracted from approximately 200 animals suspended in 50 μ l M9. RNA was extracted using Trizol (Sigma) and cDNA was synthesized using the Invitrogen Superscript Reverse Transcriptase Kit. SYBR Green Real Time PCR was carried out using the ABI 7700 and normalized to *act-1*. Primer sequences are available on request.

SUPPLEMENTAL DATA

Supplemental Data include Experimental Procedures, six tables, and fourteen figures and can be found with this article online at <http://www.cell.com/cgi/content/full/132/6/1025/DC1/>.

ACKNOWLEDGMENTS

We thank Siu Sylvia Lee, Gary Ruvkun, and Laura Mitic for generously providing advice and strains; Stacey Robida, Anne Oelmann, Jeeyong Lee, and

Michael Lucke for invaluable technical support; and members of the Blackwell lab, Javier Apfeld, and Rohit Kulkarni for critically reading this manuscript. Work was supported by funding from the NIH (grant number 2 R01 GM062891) and Iacocca Foundation (T.K.B.), the Fonds der Chemischen Industrie, BMBF FRISYS, Qualitaetsoffensive BW, and Deutsche Forschungsgemeinschaft CRC746 (R.B.), and EC 6th Framework Network of Excellence LIFE-SPAN (LSHG-CT-2007-036894) (R.B. and M.H.), as well as an NRSA (number 5 F32 GM070088-02) (R.P.O.), an NIH training grant (number T32 DK07260) (J.B.), and KRF (2006-353-C00035) and MOST/KOSEF(R112000078010010) (J.H.A.).

Received: June 29, 2007

Revised: December 18, 2007

Accepted: January 18, 2008

Published: March 20, 2008

REFERENCES

- Accili, D., and Arden, K.C. (2004). FoxOs at the crossroads of cellular metabolism, differentiation, and transformation. *Cell* 117, 421–426.
- An, J.H., and Blackwell, T.K. (2003). SKN-1 links *C. elegans* mesendodermal specification to a conserved oxidative stress response. *Genes Dev.* 17, 1882–1893.
- An, J.H., Vranas, K., Lucke, M., Inoue, H., Hisamoto, N., Matsumoto, K., and Blackwell, T.K. (2005). Regulation of the *Caenorhabditis elegans* oxidative stress defense protein SKN-1 by glycogen synthase kinase-3. *Proc. Natl. Acad. Sci. USA* 102, 16275–16280.
- Antebi, A. (2007). Genetics of aging in *Caenorhabditis elegans*. *PLoS Genet.* 3, 1565–1571. 10.1371/journal.pgen.0030129.
- Arantes-Oliveira, N., Apfeld, J., Dillin, A., and Kenyon, C. (2002). Regulation of life-span by germ-line stem cells in *Caenorhabditis elegans*. *Science* 295, 502–505.
- Berdichevsky, A., Viswanathan, M., Horvitz, H.R., and Guarente, L. (2006). *C. elegans* SIR-2.1 interacts with 14–3-3 proteins to activate DAF-16 and extend life span. *Cell* 125, 1165–1177.
- Bishop, N.A., and Guarente, L. (2007). Two neurons mediate diet-restriction-induced longevity in *C. elegans*. *Nature* 447, 545–549.
- Bowerman, B., Eaton, B.A., and Priess, J.R. (1992). *skn-1*, a maternally expressed gene required to specify the fate of ventral blastomeres in the early *C. elegans* embryo. *Cell* 68, 1061–1075.
- Brunet, A., Park, J., Tran, H., Hu, L.S., Hemmings, B.A., and Greenberg, M.E. (2001). Protein kinase SGK mediates survival signals by phosphorylating the forkhead transcription factor FKHRL1 (FOXO3a). *Mol. Cell. Biol.* 21, 952–965.
- Cahill, C.M., Tzivion, G., Nasrin, N., Ogg, S., Dore, J., Ruvkun, G., and Alexander-Bridges, M. (2001). Phosphatidylinositol 3-kinase signaling inhibits DAF-16 DNA binding and function via 14–3-3-dependent and 14–3-3-independent pathways. *J. Biol. Chem.* 276, 13402–13410.
- Dillin, A., Hsu, A.L., Arantes-Oliveira, N., Lehrer-Graiwer, J., Hsin, H., Fraser, A.G., Kamath, R.S., Ahringer, J., and Kenyon, C. (2002). Rates of behavior and aging specified by mitochondrial function during development. *Science* 298, 2398–2401.
- Finkel, T., and Holbrook, N.J. (2000). Oxidants, oxidative stress and the biology of ageing. *Nature* 408, 239–247.
- Friedman, D.B., and Johnson, T.E. (1988). A mutation in the *age-1* gene in *Caenorhabditis elegans* lengthens life and reduces hermaphrodite fertility. *Genetics* 118, 75–86.
- Gems, D., Sutton, A.J., Sundermeyer, M.L., Albert, P.S., King, K.V., Edgley, M.L., Larsen, P.L., and Riddle, D.L. (1998). Two pleiotropic classes of *daf-2* mutation affect larval arrest, adult behavior, reproduction and longevity in *Caenorhabditis elegans*. *Genetics* 150, 129–155.
- Goldstein, B.J., Mahadev, K., and Wu, X. (2005). Redox paradox: insulin action is facilitated by insulin-stimulated reactive oxygen species with multiple potential signaling targets. *Diabetes* 54, 311–321.

- Gottlieb, S., and Ruvkun, G. (1994). *daf-2*, *daf-16* and *daf-23*: genetically interacting genes controlling Dauer formation in *Caenorhabditis elegans*. *Genetics* 137, 107–120.
- Greer, E.L., Dowlatshahi, D., Banko, M.R., Villen, J., Hoang, K., Blanchard, D., Gygi, S.P., and Brunet, A. (2007). An AMPK-FOXO pathway mediates longevity induced by a novel method of dietary restriction in *C. elegans*. *Curr. Biol.* 17, 1646–1656.
- Hayes, J.D., and McMahon, M. (2001). Molecular basis for the contribution of the antioxidant responsive element to cancer chemoprevention. *Cancer Lett.* 174, 103–113.
- Henderson, S.T., and Johnson, T.E. (2001). *daf-16* integrates developmental and environmental inputs to mediate aging in the nematode *Caenorhabditis elegans*. *Curr. Biol.* 11, 1975–1980.
- Hertweck, M., Gobel, C., and Baumeister, R. (2004). *C. elegans* SGK-1 is the critical component in the Akt/PKB kinase complex to control stress response and life span. *Dev. Cell* 6, 577–588.
- Houthoofd, K., Braeckman, B.P., Lenaerts, I., Brys, K., De Vreese, A., Van Eygen, S., and Vanfleteren, J.R. (2002). Axenic growth up-regulates mass-specific metabolic rate, stress resistance, and extends life span in *Caenorhabditis elegans*. *Exp. Gerontol.* 37, 1371–1378.
- Houthoofd, K., Braeckman, B.P., Johnson, T.E., and Vanfleteren, J.R. (2003). Life extension via dietary restriction is independent of the Ins/IGF-1 signalling pathway in *Caenorhabditis elegans*. *Exp. Gerontol.* 38, 947–954.
- Hsin, H., and Kenyon, C. (1999). Signals from the reproductive system regulate the lifespan of *C. elegans*. *Nature* 399, 362–366.
- Hsu, A.L., Murphy, C.T., and Kenyon, C. (2003). Regulation of aging and age-related disease by DAF-16 and heat-shock factor. *Science* 300, 1142–1145.
- Inoue, H., Hisamoto, N., An, J.H., Oliveira, R.P., Nishida, E., Blackwell, T.K., and Matsumoto, K. (2005). The *C. elegans* p38 MAPK pathway regulates nuclear localization of the transcription factor SKN-1 in oxidative stress response. *Genes Dev.* 19, 2278–2283.
- Kell, A., Ventura, N., Kahn, N., and Johnson, T.E. (2007). Activation of SKN-1 by novel kinases in *Caenorhabditis elegans*. *Free Radic. Biol. Med.* 43, 1560–1566.
- Kenyon, C. (2005). The plasticity of aging: insights from long-lived mutants. *Cell* 120, 449–460.
- Kenyon, C., Chang, J., Gensch, E., Rudner, A., and Tabtiang, R. (1993). A *C. elegans* mutant that lives twice as long as wild type. *Nature* 366, 461–464.
- Kimura, K.D., Tissenbaum, H.A., Liu, Y., and Ruvkun, G. (1997). *daf-2*, an insulin receptor-like gene that regulates longevity and diapause in *Caenorhabditis elegans*. *Science* 277, 942–946.
- Lakowski, B., and Hekimi, S. (1996). Determination of life-span in *Caenorhabditis elegans* by four clock genes. *Science* 272, 1010–1013.
- Larsen, P.L. (1993). Aging and resistance to oxidative damage in *Caenorhabditis elegans*. *Proc. Natl. Acad. Sci. USA* 90, 8905–8909.
- Lee, S.S., Lee, R.Y., Fraser, A.G., Kamath, R.S., Ahlinger, J., and Ruvkun, G. (2003). A systematic RNAi screen identifies a critical role for mitochondria in *C. elegans* longevity. *Nat. Genet.* 33, 40–48.
- Lin, K., Hsin, H., Libina, N., and Kenyon, C. (2001). Regulation of the *Caenorhabditis elegans* longevity protein DAF-16 by insulin/IGF-1 and germline signaling. *Nat. Genet.* 28, 139–145.
- Lithgow, G.J., and Walker, G.A. (2002). Stress resistance as a determinate of *C. elegans* lifespan. *Mech. Ageing Dev.* 123, 765–771.
- McMahon, M., Itoh, K., Yamamoto, M., Chanas, S.A., Henderson, C.J., McLellan, L.I., Wolf, C.R., Cavin, C., and Hayes, J.D. (2001). The Cap'n'Collar basic leucine zipper transcription factor Nrf2 (NF-E2 p45-related factor 2) controls both constitutive and inducible expression of intestinal detoxification and glutathione biosynthetic enzymes. *Cancer Res.* 61, 3299–3307.
- Ogg, S., Paradis, S., Gottlieb, S., Patterson, G.I., Lee, L., Tissenbaum, H.A., and Ruvkun, G. (1997). The Fork head transcription factor DAF-16 transduces insulin-like metabolic and longevity signals in *C. elegans*. *Nature* 389, 994–999.
- Panowski, S.H., Wolff, S., Aguilaniu, H., Durieux, J., and Dillin, A. (2007). PHA-4/Foxa mediates diet-restriction-induced longevity of *C. elegans*. *Nature* 447, 550–555.
- Rea, S.L., Ventura, N., and Johnson, T.E. (2007). Relationship between mitochondrial electron transport chain dysfunction, development, and life extension in *Caenorhabditis elegans*. *PLoS Biol.* 5, e259. 10.1371/journal.pbio.0050259.
- Simmer, F., Tijsterman, M., Parrish, S., Koushika, S.P., Nonet, M.L., Fire, A., Ahlinger, J., and Plasterk, R.H. (2002). Loss of the putative RNA-directed RNA polymerase RRF-3 makes *C. elegans* hypersensitive to RNAi. *Curr. Biol.* 12, 1317–1319.
- Tissenbaum, H.A., and Guarente, L. (2001). Increased dosage of a *sir-2* gene extends lifespan in *Caenorhabditis elegans*. *Nature* 410, 227–230.
- Tonks, N.K. (2005). Redox redux: revisiting PTPs and the control of cell signaling. *Cell* 121, 667–670.
- van der Horst, A., and Burgering, B.M. (2007). Stressing the role of FoxO proteins in lifespan and disease. *Nat. Rev. Mol. Cell Biol.* 8, 440–450.