



Pergamon

Tetrahedron Letters 41 (2000) 6067–6069

TETRAHEDRON  
LETTERS

## An access to some functionalized azocine derivatives

Laurent Gil,<sup>a,\*</sup> Rossimiriam Pereira de Freitas Gil,<sup>b</sup> Daniela Cristina dos Santos<sup>b</sup>  
and Christian Marazano<sup>c</sup>

<sup>a</sup>*Departamento de Química, Instituto de Ciências Exatas e Biológicas, UFOP, Brazil*

<sup>b</sup>*Departamento de Química, Instituto de Ciências Exatas, UFMG, Brazil*

<sup>c</sup>*Institut de Chimie des Substances Naturelles, CNRS, 91198 Gif-sur-Yvette, France*

Received 30 March 2000; accepted 14 June 2000

---

### Abstract

The syntheses, from readily accessible 3-alkyl-4-methoxy-1,3,4,5-tetrahydropyridine **1**, of functionalized 1,6,7,8-tetrahydroazocine **7** and 1,2,7,8-tetrahydroazocine **9** are reported. © 2000 Published by Elsevier Science Ltd.

*Keywords:* azocines; nitrogen heterocycles.

---


Eight-membered nitrogen heterocycles constitute an important class of compounds, especially in view of their pharmacological properties.<sup>1</sup> These derivatives are generally difficult to obtain<sup>2</sup> and, accordingly, relatively few methods are available for their preparation. This is especially the case of highly functionalized derivatives or derivatives suitable for further chemoselective reactions.

We recently reported<sup>3</sup> that tetrahydropyridines **1** (Scheme 1, R<sup>1</sup> and R<sup>2</sup> = alkyl group) are good precursors of 3-alkyl 1,6-dihydropyridines **2** which were too unstable to be isolated, but could be trapped with common dienophiles to produce isoquinuclidine derivatives **3**. As a continuation of our studies concerning the reactivity of derivatives **1**, we now report that the reaction of these reactive intermediates with ethyl propiolate gave highly functionalized azocines heterocycles **4** in high yield.


Dihydropyridinium salt **5** (Scheme 2) was obtained from the corresponding pyridinium salt following well-established procedures.<sup>4</sup> The reaction of salt **5** with sodium methoxide in methanol gave adduct **1a** in good yield.<sup>3</sup> Adduct **1a**, when treated with ethyl propiolate in refluxing acetonitrile, gave the functionalized 1,6,7,8-tetrahydroazocine<sup>5</sup> **7** in 90% yield. It is believed that azocine formation involves an initial [2+2] cycloaddition<sup>3,6</sup> reaction between the enamine **1a** to give the cyclobutene intermediate **6**, followed by an electrocyclic ring opening to give the azocine system.

---

\* Corresponding author. Fax: +55 031 551 1707; e-mail: laurent@iceb.ufop.br



Scheme 1.



Scheme 2.

Azocine **7** was treated with 1 equivalent of methanesulfonic acid to give, presumably, an iminium salt (**8**, not isolated). Reduction of this salt with NaBH<sub>4</sub> in methanol finally gave 1,2,7,8-tetrahydroazocine **9**.

Further studies to extend this approach to other azocines are currently in progress.

## Acknowledgements

We thank FAPEMIG-MG and CAPES, Brazil for financial support.

## References

1. Evans, P. A.; Holmes, A. B. *Tetrahedron* **1991**, *47*, 9131–9166.
2. For previous related approaches, see: Lallemand, M.-C.; Chiadmi, M.; Tomas, A.; Kunesch, N.; Husson, H.-P. *Tetrahedron Lett.* **1995**, *36*, 2053–2056, and references cited therein.
3. Gil, L.; Gateau-Olesker, A.; Marazano, C.; Das, B. C. *Tetrahedron Lett.* **1995**, *36*, 707–710.
4. Grierson, D. S.; Harris, M.; Husson, H. P. *J. Am. Chem. Soc.* **1980**, *102*, 1064–1082.
5. Selected data for **7**: mp 112°C (ether–heptane); IR (cm<sup>-1</sup>) 2981, 2931, 1685, 1605, 1577, 1453, 1421, 1364, 1291, 1251, 1196; <sup>1</sup>H NMR (CDCl<sub>3</sub>, 200 MHz) δ 1.03 (1H, m), 1.29 (3H, t, *J* = 7.1 Hz), 1.57 (1H, tt, *J* = 12.6, 5 Hz), 1.64 (3H, d, *J* = 1.6, 1.8 Hz), 2.85 (1H, m), 3.24 (3H, s), 3.60 (1H, m), 4.00–4.40 (3H, m), 4.20 (2H, q, *J* = 7.1 Hz), 6.28 (1H, m), 7.23–7.40 (5H, m), 7.61 (1H, s); <sup>13</sup>C NMR (CDCl<sub>3</sub>, 50.32 MHz) δ 14.74, 16.82, 21.12, 44.98, 57.07, 59.82, 61.51, 79.21, 95.26, 122.39, 127.78, 128.18, 128.93, 131.33, 136.61, 148.99, 169.88; HRMS calcd for C<sub>19</sub>H<sub>25</sub>NO<sub>3</sub>: 315.1835; found: 315.1842; anal. calcd for C<sub>19</sub>H<sub>25</sub>NO<sub>3</sub>: C, 72.35; H, 7.99; N, 4.44; O, 15.22; found: C, 72.07; H, 8.24; N, 4.53; O, 15.03.
6. Weinstein, B.; Lin, L.-C. C.; Fowler, F. W. *J. Org. Chem.* **1980**, *45*, 1657–1661.